


## Case report

## Contained aortic rupture in a term pregnant patient during the COVID-19 pandemic

Kelly Bogaert <sup>1</sup>, Kyrstin Christensen,<sup>1</sup> Matthew Cagliostro,<sup>2</sup> Lauren Ferrara<sup>3,4</sup><sup>1</sup>Department of Obstetrics, Gynecology, and Reproductive Science, Icahn School of Medicine at Mount Sinai, New York, New York, USA<sup>2</sup>Department of Cardiology, Icahn School of Medicine at Mount Sinai, New York, New York, USA<sup>3</sup>Department of Maternal Fetal Medicine, Obstetrics and Gynecology, Icahn School of Medicine at Mount Sinai, New York, New York, USA<sup>4</sup>Department of Obstetrics and Gynecology, NYC Health and Hospitals Elmhurst, Elmhurst, New York, USA**Correspondence to**Dr Kelly Bogaert;  
kelly.bogaert@mountsinai.org

KB and KC are joint first authors.

Accepted 2 September 2020

**SUMMARY**

Aortic dissection and rupture is a rare occurrence in pregnant and postpartum patients. This case discusses the presentation and diagnosis of a patient with an acute contained thoracic aortic aneurysm rupture at 38 weeks of gestation, after presenting with throat pain and syncope during the COVID-19 pandemic. The patient underwent emergent caesarean delivery for non-reassuring fetal heart tracing, following which continued syncope workup revealed an aortic aneurysm and pericardial effusion. Diagnosis in this case was finalised with multimodality imaging, including transthoracic echocardiogram, and the patient underwent surgical aortic repair.

**BACKGROUND**

Aortic dissection and rupture are rare but life-threatening conditions, particularly during pregnancy. Most commonly, these conditions arise in the third trimester or postpartum period, with the most common cause of death being cardiac tamponade.<sup>1,2</sup> Aortic dissections are stratified into two categories based on the Stanford Criteria, with Type A involving the ascending aorta and Type B not involving the ascending aorta.<sup>3</sup> Type A aortic dissections are more common during pregnancy and postpartum, and can lead to a mortality rate as high as 50% if not treated surgically within 48 hours.<sup>4-6</sup> Risk factors for aortic dissection include connective tissue disorders, such as Marfan syndrome and Loeys-Dietz syndrome, congenital cardiovascular anomalies, such as bicuspid aortic valve and aortic




**Figure 2** Coronal view of CT angiogram of the chest, moderate pericardial effusion and no pulmonary embolus.

coarctation, familial history of aortic dissection and chronic hypertension.<sup>1,3,7</sup>

We present the case of a patient diagnosed with an acute aortic rupture at term during the COVID-19 pandemic. While previous case reports have described the diagnosis and management of aortic dissection and rupture during pregnancy, this case uniquely describes the challenges of diagnosing aortic dissection and rupture during the COVID-19 pandemic as well as the use of echocardiogram in determining this diagnosis.

**CASE PRESENTATION**

A 33-year-old female gravida 1 presented at 38 weeks of gestation to a hospital in Queens, New York, after a syncopal episode. The patient had received routine prenatal care with a private physician, and her pregnancy thus far had been uncomplicated. She reported feeling well the day prior to presentation, but woke from sleep around 04:00 on the day of presentation with shortness of breath, throat pain and 'clamminess'. The patient's husband woke to her falling off the bed, having lost consciousness and subsequently hitting her head and abdomen. On waking, the patient endorsed dizziness, mild shortness of breath and throat pain. On arrival to labour and delivery triage, the patient was noted to have oligohydramnios on pelvic ultrasound with an amniotic fluid index of 1, and fetal tachycardia with minimal to moderate variability. The patient herself was tachycardic to 109 beats/min with a blood pressure of 88/66 mm Hg, vital signs that are considered within normal limits for a pregnant patient. The patient was given a bolus of intravenous fluids, and a COVID-19 swab was sent in addition to standard inflammatory markers recommended in the work-up of COVID-19.



**Figure 1** Chest X-ray on hospital day 1 with calcification 3.5 cm below the carina (arrow).



© BMJ Publishing Group Limited 2020. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Bogaert K, Christensen K, Cagliostro M, et al. *BMJ Case Rep* 2020;**13**:e238370. doi:10.1136/bcr-2020-238370




**Figure 3** Lateral view of CT angiogram of the chest, ascending thoracic aortic aneurysm measuring 4.5 cm in anteroposterior (AP) diameter.

### INVESTIGATIONS


An ECG demonstrated sinus tachycardia with non-specific T-wave inversions in lead III. A portable chest X-ray was obtained and demonstrated a calcification 3.5 cm below the carina with some suggestion of mediastinal widening, which was not commented on in the report (figure 1).

Given the non-reassuring fetal status and lack of improvement with intrauterine resuscitation with intravenous fluids and positioning, the decision was made to proceed with caesarean delivery under spinal anaesthetic as the patient was asymptomatic and hemodynamically stable. The patient underwent an uncomplicated primary caesarean section using spinal anaesthetic, delivering a viable female infant with Apgar scores of 8 and 9 at 1 min and 5 min, respectively. After completion of the case, the patient's COVID-19 swab had returned negative. Repeat COVID-19 swab was performed given the high false-negative rate. The patient's laboratory results were otherwise significant for a troponin of 0.056 ng/mL and a D-dimer of 2301 ng/mL.

Given the patient's history of a syncopal episode with an elevated D-dimer and troponin level, there was concern for a



**Figure 4** Point of care transthoracic echocardiogram demonstrating large pericardial effusion with apparent clotted blood within pericardial space and ascending aortic root aneurysm measuring 5.2 cm without obvious dissection.



**Figure 5** Point of care transthoracic echocardiogram demonstrating large pericardial effusion with apparent clotted blood within pericardial space and ascending aortic root aneurysm measuring 5.2 cm without obvious dissection.

pulmonary embolism with right heart strain as the aetiology for her presentation. Subsequently, a CT angiogram of the chest was obtained and demonstrated moderate pericardial effusion and an ascending thoracic aortic aneurysm measuring 4.5 cm in anteroposterior (AP) diameter without dissection; no pulmonary embolus was present (figures 2 and 3).

The cardiology team was paged and performed a transthoracic echocardiogram (TTE) at bedside with findings of a large pericardial effusion and apparent clotted blood within the pericardial space, as well as an ascending aortic root aneurysm measuring 5.2 cm without obvious dissection (figures 4 and 5). The team then discovered that the patient's father had died at age 40 from an acute aortic event.

### TREATMENT

Due to concern for a concealed dissection and possible need for surgical intervention, the patient was urgently transferred to a cardiothoracic surgery team at a quaternary centre. There she underwent urgent valve-sparing aortic root replacement. Intraoperatively, the surgical team found significant hemopericardium with clotted blood, as noted on the TTE, with a ~1 cm contained ascending thoracic aortic rupture tamponaded by the pulmonary artery in the setting of a bicuspid aortic valve and ascending aortic aneurysm.

### OUTCOME AND FOLLOW-UP

The patient's post-operative course was uncomplicated from both an obstetric and cardiothoracic standpoint and she was discharged home on postoperative day 5. As an outpatient, she subsequently underwent genetic testing and was diagnosed with Loews-Dietz syndrome secondary to an SMAD3 mutation.

### DISCUSSION

Aortic dissection or rupture is a rare occurrence in pregnancy and postpartum. Aortic dissection during pregnancy makes up only 29 cases out of 9707 total cases documented by the International Registry of Acute Aortic Dissections over the past 20 years.<sup>8</sup> Aortic dissection or rupture occurred in 36 out of 6.6 million pregnancies in a study by Kamel *et al.*<sup>9</sup> Despite the rare occurrence of this disease, pregnancy itself is a risk factor for aortic pathology. In women under 40 years old, up to 50% of aortic dissections occur during pregnancy or postpartum.<sup>3</sup>

The incidence of aortic dissection or rupture is 5.5 per million patients during pregnancy and postpartum, as compared with 1.7 per million among non-pregnant patients.<sup>9</sup> In patients with aortic dissection during pregnancy, up to 62% are diagnosed with an underlying predisposing pathology.<sup>8</sup> The most commonly inherited and congenital conditions include Marfan syndrome, bicuspid aortic valve and Loeys-Dietz syndrome.<sup>8</sup> Obtaining an accurate family history can be essential in eliciting a patient's risk for this condition during pregnancy. Our patient noted history of sudden death in her father around age 40 from an aortic dissection or rupture and was subsequently diagnosed with Loeys-Dietz syndrome. Loeys-Dietz syndrome is an autosomal dominant connective-tissue disorder due to genetic mutations, such as in the SMAD3 gene, as in our patient.<sup>10</sup> It confers higher risk to patients during pregnancy, including an increased risk of aortic dissection in the third trimester.<sup>10</sup> A thorough family history not only aids in making this rare diagnosis, but is also essential for preconception counselling in these patients.

Diagnosis of aortic dissection and rupture requires reliance on history, clinical examination and imaging. While the classic presentation of aortic dissection includes sharp chest pain radiating to the back and dyspnoea, syncope can be present in up to 13% of cases.<sup>11</sup> In case reports of aortic dissection in pregnancy, symptoms have included tearing chest pain, back pain, dyspnoea, hypertension, nausea/vomiting, syncope and seizure.<sup>12–15</sup> The broad overlap between these symptoms and other more common complaints of pregnancy such as labour, pre-eclampsia and peripartum cardiomyopathy may confuse the clinician in diagnosing an aortic dissection. The gold standard for diagnosing aortic aneurysm and dissection is the CT angiogram.<sup>16</sup> If aortic disease is suspected antepartum, MRI can also be used, but the use is limited to hemodynamically stable patients.<sup>17</sup> Studies have additionally shown the utility of the echocardiogram as a tool for diagnosis given its high sensitivity and specificity.<sup>16</sup> Transesophageal echocardiogram has a higher sensitivity and specificity for diagnosing aortic dissection than TTE, but is an invasive procedure and less available in some places.<sup>17</sup> TTE is a cost-efficient diagnostic modality that may aid in diagnosing hemopericardium, increasing diagnostic concern for aortic dissection or rupture, and allow for assessment of the aortic root size and the presence of bicuspid aortic valve.<sup>17</sup> The TTE findings seen in this case, along with the patient's history of syncope, were the predominant clinical clues for the cardiology team to determine the diagnosis of aortic aneurysm rupture.

At the time of our patient's presentation, the COVID-19 pandemic both further complicated and aided in her diagnosis. Up to 37% of pregnant patients with COVID-19 have presented with dyspnoea.<sup>18</sup> During the work-up of the patient's syncope and dyspnoea, a D-dimer test was sent, a laboratory test that is frequently excluded from a dyspnoea work-up in pregnant patients given baseline elevations in D-dimer levels for healthy pregnant women.<sup>19</sup> During the COVID-19 pandemic, however, D-dimer elevation has been associated with severe pneumonia in pregnancy and is included in the standard COVID-19 panel at our hospital.<sup>18</sup> The elevated D-dimer and negative COVID-19 testing in our patient spurred further work-up for pulmonary embolism, despite her improved symptoms, and ultimately led to her diagnosis.

Aortic dissection and rupture are rare occurrences in pregnant patients, and require a high index of clinical suspicion given the similarity of symptoms with other pathologies during pregnancy, particularly during the COVID-19 pandemic. This case presents a rare pathology in a 33-year-old pregnant woman, in which

COVID-19 testing and echocardiography served to clinch the diagnosis and facilitate transfer for urgent surgical intervention.

### Learning points

- ▶ While the gold standard for diagnosing aortic aneurysm and dissection is CT angiogram, echocardiogram has the potential to quickly aid in diagnosis given its high sensitivity and specificity.
- ▶ Maintaining a broad differential is essential to patient care during the COVID-19 pandemic, as unrelated diagnoses with similar presentations to COVID-19 may still affect patients.
- ▶ In pregnancy, a patient presenting with aortic dissection may have similar symptoms to more common complaints such as labour, pre-eclampsia and peripartum cardiomyopathy.
- ▶ More than half of pregnant patients with aortic dissection in pregnancy have an underlying predisposing pathology; eliciting a thorough family history is essential to care of these women.

**Acknowledgements** We thank Drs Thomas Marino and Gaetano Bello for their instrumental role in the patient's care and clinical decisions.

**Contributors** KB and KC contributed equally as first author to the paper. KB, KC and MC all participated in the patient's care. All authors contributed to the writing of the manuscript and approved the final manuscript.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent for publication** Obtained.

**Provenance and peer review** Not commissioned; externally peer reviewed.

This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may use, download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

### ORCID iD

Kelly Bogaert <http://orcid.org/0000-0002-7376-7558>

### REFERENCES

- 1 Rajagopalan S, Nwazota N, Chandrasekhar S. Outcomes in pregnant women with acute aortic dissections: a review of the literature from 2003 to 2013. *Int J Obstet Anesth* 2014;23:348–56.
- 2 Underhill S, Woods S, Froelicher E, et al. *Cardiac nursing*. 2nd edn. Philadelphia: JB Lippincott, 1989.
- 3 Patel PD, Arora RR, Pathophysiology ARR. Pathophysiology, diagnosis, and management of aortic dissection. *Ther Adv Cardiovasc Dis* 2008;2:439–68.
- 4 Immer FF, Bansil AG, Immer-Bansil AS, et al. Aortic dissection in pregnancy: analysis of risk factors and outcome. *Ann Thorac Surg* 2003;76:309–14.
- 5 De Martino A, Morganti R, Falcetta G, et al. Acute aortic dissection and pregnancy: review and meta-analysis of incidence, presentation, and pathologic substrates. *J Card Surg* 2019;34:1591–7.
- 6 Scott C, Burruss N, Kalimi R, et al. Acute ascending aortic dissection during pregnancy. *Am J Crit Care* 2001;10:430–3.
- 7 Baue AE, Geha A, Hammond G, et al. *Glenn's Thoracic and Cardiovascular Surgery*. 5th edn. Norwalk, CT: Appelton & Lange, 1991.
- 8 Mittauer E, Harris K, Evangelista-Masip A, et al. Aortic dissection related to pregnancy: the International registry of acute aortic dissection (IRAD). *J Am Coll Cardiol* 2020;75:2213.
- 9 Kamel H, Roman MJ, Pitcher A, et al. Pregnancy and the risk of aortic dissection or rupture: a Cohort-Crossover analysis. *Circulation* 2016;134:527–33.
- 10 Metz T. Pregnancy with Loeys-Dietz: care informed by case series. *BJOG* 2019;126:1032.
- 11 Januzzi JL, Marayati F, Mehta RH, et al. Comparison of aortic dissection in patients with and without Marfan's syndrome (results from the International registry of aortic dissection). *Am J Cardiol* 2004;94:400–2.
- 12 Mandel W, Evans EW, Walford RL. Dissecting aortic aneurysm during pregnancy. *N Engl J Med* 1954;251:1059–61.

- 13 Herman JA. Dissecting aneurysm of the ascending aorta during the third trimester of pregnancy with cardiac tamponade. *Am J Obstet Gynecol* 1965;93:282–4.
- 14 Weissmann-Brenner A, Schoen R, Divon MY. Aortic dissection in pregnancy. *Obstet Gynecol* 2004;103:1110–3.
- 15 Kinney-Ham L, Nguyen HB, Steele R, *et al*. Acute aortic dissection in third trimester pregnancy without risk factors. *West J Emerg Med* 2011;12:571–4.
- 16 Sobczyk D, Nycz K. Feasibility and accuracy of bedside transthoracic echocardiography in diagnosis of acute proximal aortic dissection. *Cardiovasc Ultrasound* 2015;13:15.
- 17 Hiratzka LF, Bakris GL, Beckman JA, *et al*. 2010 ACCF/AHA/AATS/ACR/ASA/SCA/SCAI/SIR/STS/SVM guidelines for the diagnosis and management of patients with thoracic aortic disease: a report of the American College of cardiology Foundation/American heart association Task force on practice guidelines, American association for thoracic surgery, American College of radiology, American stroke association, society of cardiovascular Anesthesiologists, Society for cardiovascular angiography and interventions, society of interventional radiology, society of thoracic surgeons, and Society for vascular medicine. *Circulation* 2010;121:e266–369.
- 18 Pereira A, Cruz-Melguizo S, Adrien M, *et al*. Clinical course of coronavirus disease-2019 in pregnancy. *Acta Obstet Gynecol Scand* 2020;99:839–47.
- 19 Hedengran KK, Andersen MR, Stender S, *et al*. Large D-dimer fluctuation in normal pregnancy: a longitudinal cohort study of 4,117 samples from 714 healthy Danish women. *Obstet Gynecol Int* 2016;2016:1–7.

Copyright 2020 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>  
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

### Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at [support@bmj.com](mailto:support@bmj.com).

Visit [casereports.bmj.com](http://casereports.bmj.com) for more articles like this and to become a Fellow